

Mammography Additional Views and Misc. Information

- **AV's**
 - Ca++:
 - Full ML
 - CC and ML Mags
 - All other abnormalities, follow what the report is asking for

- **Lumpectomy**
 - Diagnostic x 3 years - Yearly after first visit
 - 1st imaging after surgery:
 - Full CC and MLO
 - CC and ML mag
 - ML mag only each year after
 - After 3 years, patient can request a screening mammo

- **STFU:**
 - Ca++ / Post Biopsy for Ca++:
 - 6 month
 - Full CC & MLO
 - CC & ML mag
 - 1 year
 - Bilateral Mammo - Full CC and MLO's
 - CC and ML mag on ca++ being followed
 - 1.5 year / 2 year
 - Full CC and MLO's
 - CC and ML mag on ca++ being followed

 - Asymmetry / Post Biopsy for Asymmetry:
 - 6 month
 - Full CC and MLO
 - 1 year
 - Bilateral Mammo - Full CC and MLO
 - 2 year
 - Full CC and MLO's

 - Post Biopsy Distorsion
 - 3D CC & ML

 - Post Biopsy Calcs ++
 - 2D CC & ML

 - Post Biopsy Asymmetry
 - 2D CC / ML

 - Post MRI Biopsy
 - 2D Full CC & ML

 - Post Wireless Tag Placement Axilla
 - 3D Tomo for AT view

- Post Biopsy Mass
 - 2D CC / ML
-

- Nipple Discharge
 - Due for Bilateral / Unilateral:
 - Bilateral Mammo
 - Test discharge if applicable

- ADH (Atypical Ductal Hyperplasia) / ALH (Atypical Lobular Hyperplasia)
 - Non-malignant, but patient had excisional biopsy
 - Bilateral Mammo - NO MAGS

- Male Patients:
 - 25 years and older: Bilateral Mammogram
 - 24 years and younger: Ask the Radiologist for direction

- Female Patients:
 - 30 and older: Bilateral Mammogram
 - 29 and younger: Ultrasound

- Post Biopsy Mammogram:
 - Ca++, Asymmetry, Distortion, Mass
 - Full CC and ML
 - Ca++ - Mag the way they were targeted in biopsy